

A COMPREHENSIVE REVIEW OF BAMBOO LEAVES: PHYTOCHEMICALS, BIOACTIVITIES, TOXICITY, RESOURCE POTENTIAL IN VIETNAM

Quynh Anh Le^{1,2,3}, Thanh Tai Duong¹, Thu Thuy Thi Nguyen¹,
Ngoc Han Cao Nguyen¹, Lan Phi Thi Nguyen^{2,3*}

¹*Ho Chi Minh City University of Industry and Trade*

²*Ho Chi Minh City University of Technology (HCMUT)*

³*Vietnam National University in Ho Chi Minh City (VNU HCM)*

*Email: lanphi@hcmut.edu.vn

Received: 2 May 2025; Revised: 25 May 2025; Accepted: 1 June 2025

ABSTRACT

Bamboo leaves (*Folium bambusae*) have long been recognized as a traditional medicinal material in Eastern medicine and are extensively utilized in various classical and empirical herbal prescriptions for their heat-clearing, diuretic, and anti-inflammatory properties. In recent years, bamboo leaves have attracted considerable scientific interest, leading to an increasing number of studies. Over the past two decades, research findings related to bamboo species identification, chemical composition, biological activities, and toxicity have been consistently published. This review article synthesizes and summarizes scientific studies on the chemical composition, bioactivities, and toxicity of bamboo leaf extracts. Relevant literature was collected from international scientific databases such as PubMed, ScienceDirect, Google Scholar, Web of Science, Scopus, and ScienceGate, as well as selected reliable online sources. In addition to reviewing bamboo classification systems and global bamboo resources, this article places special emphasis on the current status and development potential of bamboo resources in Vietnam, and explores the relevance of bamboo leaves in the food industry. A total of 86 references were carefully selected based on their relevance and scientific quality and were included for data extraction and synthesis. This article provides comprehensive information on bamboo resources and various research aspects related to bamboo leaves, aiming to guide more in-depth and holistic investigations, as well as broader application trials. These efforts are expected to enhance the value of bamboo, support the sustainable development of bamboo-based ecosystems, and contribute to a green future.

Keywords: Bamboo leaves, phytochemical, bioactivities, toxicity, bamboo resources.

1. INTRODUCTION

Bamboo belongs to the grass family (Poaceae), subfamily *Bambusoideae*. It is widely distributed across tropical and subtropical regions, particularly in Asia, including China, India, Japan, and Southeast Asian countries [1]. In addition to the well-known economic values derived from bamboo culms and shoots, bamboo leaves have long been recognized as a valuable traditional medicinal material in China, India, and Japan [2]. In Vietnam, bamboo leaves are considered a traditional herbal remedy with therapeutic effects, as documented in ancient Eastern medicine texts such as “Miscellaneous Records of Famous Physicians” and “Collected Works of Tue Tinh”.

In recent years, numerous studies have been conducted to optimize the extraction processes of bamboo leaf extracts, analyze their bioactive components, and assess their toxicity. Several review articles have also been published; however, notable limitations remain. For instance, the review titled “Miraculous medicinal properties of Bamboo: a Review” by Indian authors (2021) describes various medicinal properties of bamboo but lacks reliable citations and does not comprehensively cover scientific evidence on the phytochemical composition and bioactivities of bamboo leaves [3]. Another review by Benjamin, M.A.Z. *et al.* (2023) focused on the *Bambusa* genus and provided detailed information on the compounds and biological activities; however, it did not specifically address bamboo leaves nor include toxicity-related studies [1]. Similarly, a review titled “Bamboo leaf: A review of traditional medicinal property, phytochemistry, pharmacology, and purification technology” published the same year offers a relatively complete overview of the phytochemical and medicinal properties of bamboo leaves but fails to summarize toxicity studies and does not discuss bamboo resource systems [2].

Therefore, this review aims to provide a comprehensive synthesis of published studies on the phytochemical composition, biological activities, and toxicity of bamboo leaves. Additionally, it compiles information on global bamboo resources, with particular emphasis on those in Vietnam, in order to guide future in-depth research and promote efficient and sustainable applications.

2. MATERIALS AND METHODS

This review was conducted using a semi-systematic approach, combining keyword-based searches with backward citation tracking from relevant review articles. The literature collection focused on four core areas:

- Phytochemical composition
- Biological activities
- Toxicity assessment
- Bamboo resources and applications

Scientific data were retrieved from major international databases including PubMed, ScienceDirect, Google Scholar, Web of Science, Scopus, SpringerLink, and ScienceGate. Additional information was collected from reputable internet-based sources such as the Food and Agriculture Organization (FAO), International Network for Bamboo and Rattan (INBAR), World Agroforestry Centre (ICRAF), Business Wire, Guadua Bamboo, and Vietnamese platforms including VNScience, Khoa học Phổ thông (a Vietnamese science outreach magazine), and the Department of Science and Technology – Ministry of Natural Resources and Environment.

The search strategy applied chronological filters from the most recent to older publications. The inclusion criteria focused on ensuring scientific relevance and quality. Specifically, documents were selected based on: (1) direct relevance to the review topic; (2) publication in reputable, peer-reviewed journals; (3) high citation frequency; and (4) availability of full-text access. Only documents meeting these criteria were included. In total, the review incorporated:

- 50 articles on phytochemical composition (published between 2009 and 2025);
- 33 articles on biological activities (published between 2013 and 2025);
- 15 articles on toxicity (published between 2006 and 2025);

- 31 sources on bamboo resources and applications include 22 scientific articles (published between 2003 and 2023) and 9 online references (updated between 2019 and 2023).

The structured process for literature identification and selection is illustrated in Figure 1.




Figure 1. Literature screening and selection process

3. RESULTS AND DISCUSSION

3.1. Phytochemical

3.1.1. Phenolic

Phenolics are naturally occurring compounds that contain one or more hydroxyl (-OH) groups directly attached to a benzene ring. These include phenolic acids, flavonoids, stilbenes, tannins, and lignans, all of which play important roles in plant growth and defense. Polyphenols are widely found in plants and are known for their beneficial biological activities in human health [4]. As a result, this group has received the most attention in bamboo leaf studies. Total phenolic content (TPC) and total flavonoid content (TFC) have been analyzed by various researchers across different bamboo species and countries, using diverse extraction conditions and methods [5–15].

The main phenolic compounds reported include:

- **Phenolic acids:** protocatechuic acid, chlorogenic acid, caffeic acid, coumaric acid, ferulic acid, 4-Hydroxybenzoic acid, and p-Coumaric acid are commonly found in most bamboo leaf species [16–29]. These acids exhibit antioxidant, anti-inflammatory, and hepatoprotective properties, and inhibit reactive oxygen species (ROS) production. Chlorogenic acid and p-Coumaric acid are considered key contributors to antioxidant activity. 3-O-caffeoyl-1-methylquinic acid, isolated from

Phyllostachys edulis (China), has demonstrated antioxidant effects, activation of cellular defense systems, and skin cancer prevention potential [25].

- **Flavonoids:** These are the most abundant and significant group of phenolics in bamboo leaves. Identified flavonoids include Isoorientin, Orientin, Homoorientin, Vitexin, Homovitexin, Isovitexin, and Luteolin. Other components include Rutin, Tricin, Apigenin, Kaempferol, Quercetin, Catechin, and Naringin (Naringenin 7-Rhamnoglucoside) [16–35]. Syringaresinol- β -D-glucoside has been identified as a marker compound for *Phyllostachys bambusoides* and *Pseudosasa japonica* (Korea) [36]. 7-O-Methyltricin was reported in *Phyllostachys edulis* [25]. Multiple studies consistently affirm that flavonoids are the most bioactive and promising compounds in bamboo leaves. Isoorientin, in particular, is recognized for its anti-inflammatory and wound-healing effects.

3.1.2. Volatile organic compounds (VOCs)

The characteristic aroma of bamboo leaves mainly derives from aldehydes, alcohols, volatile fatty acids, and furans. Representative VOCs in bamboo leaf extracts include esters, aldehydes, alcohols, β -amyrene, α -amyryn acetate, friedelin, (e)-2-hexenal, cis-3-hexen-1-ol, benzaldehyde, and octanal [7, 28, 37–39]. These compounds provide a natural pleasant fragrance and also possess antioxidant, antibacterial, antifungal, and anti-inflammatory properties. Notably, cis-3-hexen-1-ol (27–35%) was found in *Phyllostachys heterocyclus* [38] and limonene was reported in *Phyllostachys bambusoides* and *Phyllostachys pubescens*, comprising up to 32% of steam-distilled essential oil [40]. Coumaric lactone was also reported in *Phyllostachys nigra* var. *henonis* and *Phyllostachys* Sieb. et Zucc. [24, 26].

3.1.3. Alkaloids

Alkaloids have been detected in bamboo leaf extracts [11, 13, 19, 41]. The ethanol extract of *Bambusa vulgaris* contained the highest alkaloid content at 4.1% among identified compounds [5]. Alkaloids were associated with antifungal activity in studies on *Bambusa blumeana* [11].

3.1.4. Saponins

Saponins were identified in several studies. They were present in aqueous extracts of *Bambusa vulgaris* leaves and contributed to antibacterial activity [41], as well as to beneficial biochemical changes in the liver and kidneys [5].

3.1.5. Steroids and phytoestrogens

Steroids, particularly in the unsaturated form, were identified in *Bambusa vulgaris* [41]. β -Sitosterol and friedelin (phytosterols or triterpenoids) were reported in *Sasa palmata* [7]. Phytoestrogens were also found in extracts of *Bambusa bambos* (L.) Voss, primarily β -sitosterol and stigmasterol. The hydroalcoholic extract of *Bambusa bambos* has been suggested as a safe alternative to hormone replacement therapy for managing menopausal symptoms [42].

3.1.6. Other bioactive compounds

Minor compounds identified in a few studies include anthraquinones (0.06% in *Bambusa vulgaris* extract), which contribute to general bioactivity [5], and cyclitol detected via LCMS in *Bambusa stenostachya*, a naturally bioactive molecule [27].

The phytochemical profile of bamboo leaf extracts is influenced by multiple factors. Various extraction methods have been employed across studies, ranging from conventional techniques such as Soxhlet extraction and maceration [7, 15], to modern approaches including pressurized liquid extraction, microwave-assisted extraction, and ultrasound-assisted extraction [8, 10, 23, 32, 43]. A common limitation in these investigations is the insufficient assessment of influencing parameters, as most studies are conducted under a single condition or involve limited optimization, thereby restricting the broader applicability of the results. The age of the bamboo plant and the timing of leaf collection have been shown to significantly impact both the phytochemical content and the bioactivity of the extracts [14, 23]. However, very few studies have addressed the effects of external factors such as environmental conditions, soil type, or leaf maturity, which are likely to contribute meaningfully to phytochemical variability. Advanced analytical techniques such as HPLC, RP-HPLC, UHPLC, LC-MS/MS, and UPLC-QTOF-MS/MS have been widely used to identify specific phytochemicals in bamboo leaf extracts [13, 16, 17, 22, 23, 30, 31, 42, 44]. The current literature largely focuses on well-established compound classes, including phenolic acids, flavonoids, and volatile compounds. A broader exploration of less-characterized phytochemical groups is needed to achieve a more comprehensive understanding of the phytochemical complexity of bamboo leaves.

3.2. Biological activities

3.2.1. Antioxidant activity

Antioxidant activity is one of the most characteristic properties of phytochemicals, particularly phenolic compounds such as flavonoids [4]. The antioxidant potential of bamboo leaf extracts has been demonstrated in numerous studies and across various bamboo species [6, 9, 10, 13-17, 19-23, 26, 28, 34, 38, 40, 43, 45-49]. Dried leaves from *Bambusa* spp., *Dendrocalamus* spp., and *Thyrsostachys siamensis* have shown strong antioxidant effects - comparable to green tea and superior to many other herbs such as hibiscus or lemongrass [12] (Table 1).

3.2.2. Antibacterial and antifungal activities

Different bamboo species have demonstrated broad-spectrum antibacterial and antifungal effects, particularly *Phyllostachys pubescens*, *Bambusa blumeana*, and *Bambusa vulgaris*. Extracts from *Bambusa* spp., *Phyllostachys* spp., *Dendrocalamus strictus*, and *Chimonocalamus delicatus* have been shown to inhibit several bacterial strains, including *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Escherichia coli*, *Staphylococcus epidermidis*, and *Klebsiella pneumoniae* [15, 19, 22, 38, 46, 48]. Phenolic-rich extracts from *Bambusa vulgaris* leaves exhibited strong antibacterial effects [50]. At the 12th World Bamboo Congress (2024), it was affirmed that flavonoid-rich bamboo leaf extracts are effective against a wide range of bacteria [51].

Fungal strains inhibited by bamboo extracts include *Phytophthora capsici*, *Fusarium graminearum*, *Valsa mali*, *Botryosphaeria dothidea*, *Venturia nashicola*, *Botrytis cinerea*, *Colletotrichum musae*, *Alternaria alternata*, *Aspergillus niger*, and *Penicillium chrysogenum* [11, 14, 19, 52] (Table 1).

3.2.3. Anticancer activity

In vitro studies on breast and prostate cancer cells showed that extracts from *Bambusa arundinacea*, *Phyllostachys* spp., *Guadua incana* possess anticancer properties [40, 48, 53] (Table 1).

3.2.4. Anti-inflammatory and vasoprotective effects

Extracts of *Phyllostachys edulis* reduced the expression of pro-inflammatory cytokines, likely due to phenolic and flavonoid compounds such as isoorientin, orientin, and isovitexin [16]. Both orientin and isoorientin inhibited cytokine activity and exhibited vasoprotective effects by suppressing lipopolysaccharide (LPS)-induced inflammation [54] (Table 1).

3.2.5. Anti-hyperglycemic activity

Hydroalcoholic extracts from *Bambusa bambos* leaves significantly reduced blood glucose levels in streptozotocin (STZ)-induced diabetic mice. At a dose of 400 mg/kg, the effect was comparable to the standard drug metformin [55] (Table 1).

3.2.6. Neuroprotective activity

In Alzheimer's disease model mice, bamboo leaf extracts were shown to regulate neurotransmitter pathways, inhibit neurotransmitter-degrading enzymes, and improve learning and memory function, demonstrating neuroprotective effects [56] (Table 1).

3.2.7. Anti-allergic effects

Extracts from *Phyllostachys pubescens* leaves significantly inhibited IgE production in peripheral blood lymphocytes, reducing levels by up to 42.8% compared to the control, indicating strong anti-allergic potential [22] (Table 1).

Table 1. Biological activities of bamboo leaves

No.	Biological Activity	Species	Countries	References
1	Antioxidant	<i>Bambusa</i> spp., <i>Dendrocalamopsis</i> spp., <i>Dinochloa</i> spp., <i>Gigantochloa</i> spp., <i>Lophatherum</i> spp., <i>Pleioblastus</i> spp., <i>Phyllostachys</i> spp., <i>Chimonocalamus delicatus</i> , <i>Shibataea</i> spp., <i>Thyrsostachys</i> spp., <i>Schizostachyum</i> spp.	India, Malaysia, Myanmar, Nepal, Japan, China, Korea	[4, 6, 9, 10, 13-17, 19-23, 26, 28, 34, 38, 40, 43, 45-49]
2	Antibacterial	<i>Bambusa</i> spp., <i>Phyllostachys</i> spp., <i>Chimonocalamus delicatus</i>	Myanmar, Nepal, Japan, China, India	[15, 19, 22, 38, 46, 48, 50, 51]
3	Antifungal	<i>Bambusa</i> spp., <i>Phyllostachys</i> spp.	Philippines, China	[11, 14, 19, 52]
4	Anticancer	<i>Bambusa</i> spp., <i>Phyllostachys</i> spp.	India, Korea	[40, 48, 53]
5	Anti-inflammatory, Vasoprotective	<i>Phyllostachys edulis</i> J. Houz	China	[16, 54]

6	Anti-hyperglycemic	<i>Bambusa bambos</i>	India	[55]
7	Neuroprotective	Bamboo leaf extract	China	[56]
8	Anti-allergic	<i>Phyllostachys pubescens</i>	Japan	[22]

With a diverse range of phytochemicals, bamboo leaf extracts have demonstrated a wide spectrum of biological activities across various studies. However, most findings are limited to in vitro experiments conducted at a few selected concentrations, often without appropriate control groups for comprehensive efficacy evaluation. Future research should delve deeper into the underlying biological and chemical mechanisms and expand toward in vivo investigations to validate and broaden current findings.

3.3. Toxicity evaluation

To ensure the safe application of bamboo leaves in food and health products, toxicity testing is essential. *In vitro* studies using extracts of *Phyllostachys* spp. on mouse cells have shown no cytotoxic effects [27, 33, 57]. Similarly, *in vivo* studies on various organisms such as *Artemia salina* larvae, mice, and rabbits have demonstrated the safety of bamboo leaf extracts from *Bambusa* spp., *Phyllostachys* spp., *Dinochloa sublaevigata*, *Gigantochloa levis*, and *Schizostachyum brachycladum* [5, 13, 24, 25, 35, 55, 58]. Dietary supplementation with bamboo leaf powder or extract in broiler chickens did not adversely affect physiological parameters and even improved growth performance and oxidative stress resistance [59,60] (Table 2).

Recently, six compounds (p-Coumaric acid, ferulic acid, caffeic acid, protocatechuic acid, gallic acid, and vanillic acid) were evaluated through in silico methods by Malaysian researchers and were found to be non-toxic and suitable for potential development into antimicrobial drugs [50] (Table 2).

However, some studies have indicated that extracts from *Phyllostachys edulis* may cause cytotoxicity in human liver cells (HepG2) at high concentrations, although they are safe at lower levels [16, 18] (Table 2).

In summary, bamboo leaves appear safe in preclinical assessments, but current toxicity studies remain limited to a few species and are primarily conducted in vitro or in animal models. Therefore, further toxicity testing, including clinical trials, is needed to provide a more comprehensive safety profile and to support the wider application of bamboo leaves.

Table 2. Toxicological studies on bamboo leaves

No.	Test Type	Bamboo Species	Results	Reference
1	In vitro on HaCaT and 3T3 fibroblasts	<i>Phyllostachys edulis</i>	Non-toxic at 10–250 µg/mL and 5–100 µM	[33]
2	In vitro on HepG2	<i>Phyllostachys edulis</i>	Cytotoxic at 0.1 and 0.2 mg/mL	[16]
3	In vitro on HepG2	<i>Phyllostachys pubescens</i>	Safe at ≤120 µg/mL; toxic at >240 µg/mL	[18]
4	In vitro on mouse microglia (BV2 cells)	<i>Phyllostachys</i> spp. (Isoorientin)	Safe at 20 µM; neuroprotective	[57]

No.	Test Type	Bamboo Species	Results	Reference
5	In vitro on NIH/3T3 cells	<i>Bambusa stenostachya</i>	Non-toxic per ISO standard	[27]
6	In vivo on Artemia salina larvae	<i>Bambusa</i> spp., <i>Dinochloa sublaevigata</i> , <i>Gigantochloa levis</i> , <i>Schizostachyum brachycladum</i>	All samples non-toxic (LC ₅₀ > 1000 µg/mL)	[13]
7	In vivo on ICR and Sprague Dawley mice	<i>Phyllostachys nigra</i> var. <i>henonis</i>	Safe at ≤4.3 g/kg/day; no toxicity	[27]
8	In vivo on Swiss albino mice	<i>Bambusa bambos</i>	No acute toxicity at 2000 mg/kg	[55]
9	In vivo on Sprague Dawley mice	<i>Phyllostachys nigra</i> var. <i>henonis</i> Strapf	No toxicity at <5000 mg/kg	[58]
10	In vivo on Sprague–Dawley & Kunming mice	<i>Phyllostachys nigra</i> var. <i>henonis</i>	Safe up to 4.3 g/kg/day	[24, 35]
11	In vivo on pregnant Dutch rabbits	<i>Bambusa vulgaris</i>	No histological damage, but affected liver, kidney, and hormones	[5]
12	In vivo on broiler chickens	<i>Bambusa vulgaris</i>	Reduced heat stress, improved growth at 1–2%	[59]
13	In vivo on broiler chickens	<i>Bambusa vulgaris</i>	Improved feed conversion, growth, reduced oxidative stress	[60]
14	In silico prediction	<i>Bambusa vulgaris</i>	No predicted toxicity	[50]

Toxicological studies on bamboo leaf extracts remain limited in terms of species coverage and published data. Current findings are largely confined to in vivo experiments in animals, conducted at specific dosage levels. Future research should focus on clinical trials in both animals and humans to advance the application potential of bamboo leaf extracts, particularly in the food industry.

3.4. Global and Vietnamese bamboo resources

According to data from the Royal Botanic Gardens, Kew (updated as of July 2023), bamboo belongs to the grass family (Poaceae), subfamily *Bambusoideae*, and is categorized into three main tribes:

Bambuseae (tropical woody bamboos, the most common),

Arundinarieae (temperate bamboos), and

Olyreae (herbaceous bamboos that typically grow under forest canopies).

Each tribe contains numerous genera and species, totaling 139 genera and approximately 1,821 identified species worldwide [61]. The most commonly cultivated species is *Bambusa vulgaris* [62] (Figure 2).




Figure 2. Taxonomic classification of bamboo (Bambusoideae)

The global bamboo coverage is estimated at approximately 35 million hectares, representing around 1% of the world's forest area [63, 64]. Although taxonomically a grass, bamboo offers ecological and economic benefits comparable to, and in some cases exceeding, those of industrial tree species such as *Acacia* and *Eucalyptus*. Its principal advantages include a rapid growth rate, high planting density, and significant biomass yield. Additionally, its short harvesting cycle enhances its economic viability [63-66]. The global bamboo market is projected to grow at a compound annual growth rate (CAGR) of 5.7% from 2021 to 2028, reaching an estimated value of USD 82.9 billion by 2028 [67].

Bamboo is primarily distributed in Asia, Africa, and the Americas, with Asia accounting for approximately 55% of the global bamboo area. Major bamboo-producing countries include China, India, Vietnam, and Japan [1-3, 64, 68–70] (Figure 3).




Figure 3. Bamboo distribution by region

Bamboo Resources in Vietnam

Vietnam’s tropical monsoon climate is highly favorable for bamboo growth. The country has around 1.6 million hectares of bamboo, found across all ecological regions—from mountains and plains to coastal and island areas. Among Vietnam’s 63 provinces, 37 have more than 10,000 hectares of bamboo [71, 72]. Bamboo accounts for about 15% of Vietnam’s natural forest area, and notably, 95% of bamboo coverage in the country is in natural forests, primarily located in the Northern Midlands and Mountains, North Central, and Central Highlands regions [71, 73] (Figure 4).




Figure 4. Bamboo Area by Province (Natural vs. Total)

Vietnam is home to approximately 26 genera and 216 species of bamboo, with the most diverse genera being *Bambusa* (60–70 species), *Dendrocalamus* (20–25 species), *Gigantochloa* (10–16 species), *Schizostachyum* (14 species), and *Indosasa* (11 species) [74, 75].

Common and widely distributed species in Vietnam include: *Bambusa blumeana* (thorny bamboo), *Bambusa vulgaris*, *Schizostachyum* spp., *Gigantochloa* spp. [70, 76].

Most of these species are native, while 16.7% are introduced [62]. Endemic or recently identified species include *Chimonocalamus bidoupensis*, *Yersinochloa nghiana*, *Yersinochloa dalatensis*, *Cochinchinochloa braiana*, *Annamocalamus kontumensis*, and *Nianhochloa bidoupensis* [77–81].

These findings underscore bamboo's crucial role in conserving Vietnam's biodiversity and contributing to global ecological sustainability. An article published by the Khoa học Phổ thông magazine (under the HCMC People's Committee) emphasized bamboo's potential for forest restoration and climate change mitigation. Vietnamese bamboo is recognized as a "champion species" for its environmental benefits. It grows rapidly, generates large biomass in a short time, and sequesters significantly more carbon 7-30% more than fast-growing trees such as fir and eucalyptus. Additionally, bamboo roots help reduce soil erosion by up to 75%. In Vietnam, bamboo is considered a "lifesaver" in flood-prone areas. Therefore, bamboo plays a vital role in both improving livelihoods and addressing global climate challenges [82, 83].

However, Vietnam's bamboo industry remains underdeveloped. Although the country has approximately 25% of China's bamboo area, its bamboo exports are only valued at USD 300–400 million per year, while China earns up to USD 500 billion annually [84]. While the global bamboo area increased by 50% between 1990 and 2020, largely due to rapid expansion in China and India, Vietnam's bamboo area has not significantly increased and may even have declined [83]. Thus, conservation and sustainable development of Vietnam's bamboo resources is an urgent priority.

In summary, bamboo leaves are a rich source of bioactive compounds, yet they remain underutilized and largely overlooked in practical applications. In many regions, they are still mainly used as animal feed [59, 60, 85] or for wrapping food products such as cakes and traditional dishes, or for decorative purposes. As such, they are often regarded as by-products or waste materials of the bamboo industry.

Since the mid-20th century, bamboo leaves have garnered increasing scientific interest, particularly in the analysis of their chemical composition. A review of scientific publications from 2003 to the present reveals that more than 18 genera and 67 species from countries such as India, Taiwan, South Korea, Malaysia, Myanmar, Nepal, Nigeria, Japan, the Philippines, China, and Italy have been investigated for their phytochemical contents and biological activities. Applications of bamboo leaves in the food and pharmaceutical industries have also been studied and are expanding, with emerging products such as bamboo beer, bamboo leaf tea, and tea bags.

However, no scientific studies on bamboo leaves have yet been reported from Vietnam, highlighting a significant untapped potential for research and development. Promoting the use of bamboo leaves not only enhances the economic value of bamboo as a crop but also supports the conservation and sustainable development of bamboo ecosystems. Utsaphong Uprarawanna *et al.* published a comprehensive review of the nutritional and functional value of bamboo, emphasizing that bamboo leaves, not just bamboo shoots, have potential applications in food and medicine. The authors advocate for the development of value-added products from the entire bamboo plant as a viable solution to address global food security challenges [86].

4. CONCLUSION

Bamboo leaves contain a diverse array of phytochemical compounds and exhibit various biological activities, including antioxidant, antibacterial, and anti-inflammatory effects. Despite their significant potential, bamboo leaves remain largely underutilized - particularly in Vietnam, where no scientific studies have yet been published. This review highlights the potential of bamboo leaves for development in the food and pharmaceutical industries and opens new directions for research, especially in toxicity assessment and broader food-related applications. Utilizing this untapped natural resource could enhance the economic value of bamboo and promote the sustainable development of bamboo-based ecosystems.

Acknowledgement

We would like to thank Ho Chi Minh City University of Industry and Trade (HUIT); Ho Chi Minh City University of Technology (HCMUT), VNU HCM for supporting this study.

REFERENCES

1. Benjamin, M.A.Z., Saikim, F.H., Ng, S.Y., Rusdi, N.A. – A comprehensive review of the ethnobotanical, phytochemical, and pharmacological properties of the genus *Bambusa*, *J. Appl. Pharm. Sci.* 13 (5) (2023) 001–021. <https://doi.org/10.7324/JAPS.2023.98082>
2. Cheng, Y., Wan, S., Yao, L., Lin, D., Wu, T., Chen, Y., Zhang, A., Lu, C. – Bamboo leaf: A review of traditional medicinal property, phytochemistry, pharmacology, and purification technology, *J. Ethnopharmacol* 306 (2023) 116166. <https://doi.org/10.1016/j.jep.2023.116166>
3. Das, B., Singh, P.K., Munna, J. – Miraculous medicinal properties of Bamboo: a review, *J. Med. Plants Stud* 9 (5) (2021) 28–30
4. Shahidi, F., Ambigaipalan, P. – Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects – A review, *J. Funct. Foods* 18 (2015) 820–897. <https://doi.org/10.1016/j.jff.2015.06.018>
5. Yakubu, M.T., Bukoye, B.B., Oladiji, A.T., Akanji, M.A. – Toxicological implications of aqueous extract of *Bambusa vulgaris* leaves in pregnant Dutch rabbits, *Hum. Exp. Toxicol* 28 (9) (2009) 591–598. <https://doi.org/10.1177/0960327109106975>
6. Macwan, C., Patel, H.V., Kalia, K. – A comparative evaluation of in vitro antioxidant properties of bamboo *Bambusa arundinacea* leaves extracts, *J. Cell Tissue Res.* 10 (3) (2010) 2413–2418
7. Zulkafli, Z.D., Wang, H., Miyashita, F., Utsumi, N., Tamura, K. – Cosolvent-modified supercritical carbon dioxide extraction of phenolic compounds from bamboo leaves (*Sasa palmata*), *J. Supercrit. Fluids* 94 (2014) 123–129. <https://doi.org/10.1016/j.supflu.2014.07.008>
8. Shang, Y.F., Kim, S.M., Um, B.-H. – Optimisation of pressurised liquid extraction of antioxidants from black bamboo leaves, *Food Chem.* 154 (2014) 164–170. <https://doi.org/10.1016/j.foodchem.2013.12.050>
9. Narzary, R., Wary, R.R., Basumatary, J., Kalita, P., Middha, S.K., Usha, T., Goyal, A.K. – Unveiling the nutritional and bioactive potential of bamboo (*Bambusa tulda* Roxb.) tea: A phytochemical and spectroscopic study, *Adv. Bamboo Sci.* 10 (2025) 100132. <https://doi.org/10.1016/j.bamboo.2025.100132>

10. Xu, J., Wu, J., Qi, J., Li, J., Liu, Y., Miao, Z., Qiu, G., Jia, W. – Microwave-assisted extraction of flavonoids from *Phyllostachys heterocycla* leaves: Optimization, mechanism, and antioxidant activity *in vitro*, *BioResources* 16 (4) (2021) 8060–8081
11. Saduccos, A.G. – Antimycotic potential of *Kawayang tinik* against pathogenic fungal species, *Plant Sci. Today* 8 (2) (2021) 403–409. <https://doi.org/10.14719/pst.2021.8.2.1093>
12. Indira, A., Santosh, O., Koul, A., Nirmala, C. – Comparative assessment of the antioxidant potential of bamboo leaves, along with some locally and commercially consumed beverages in India, *Adv. Bamboo Sci.* 1 (2022) 100007.
13. Benjamin, M.A.Z., Ng, S.Y., Saikim, F.H., Rusdi, N.A. – The effects of drying techniques on phytochemical contents and biological activities on selected bamboo leaves, *Molecules* 27 (2022) 6458. <https://doi.org/10.3390/molecules27196458>
14. Shen, H., Wang, Y., Shi, P., Li, H., Chen, Y., Hu, T., Yu, Y., Wang, J., Yang, F., Luo, H., Yu, L. – Effects of the species and growth stage on the antioxidant and antifungal capacities, polyphenol contents, and volatile profiles of bamboo leaves, *Foods* 13 (2024) 480. <https://doi.org/10.3390/foods13030480>
15. KC, B.M., Gauchan, D.P., Khanal, S.N., Lamichhane, J. – Phytochemical screening, antioxidant and antibacterial activity of bamboo leaf collected from agroecosystem of the Central Siwalik region, Nepal, *Vegetos* (2023). <https://doi.org/10.1007/s42535-023-00761-8>
16. Tundis, R., Augimeri, G., Vivacqua, A., Romeo, R., Sicari, V., Bonofiglio, D., Loizzo, M.R. – Anti-inflammatory and antioxidant effects of leaves and sheath from bamboo (*Phyllostachys edulis* J. Houz), *Antioxidants* 12 (2023) 1239. <https://doi.org/10.3390/antiox12061239>
17. Ma, N.-H., Guo, J., Xu Chen, S.-H., Yuan, X.-R., Zhang, T., Ding, Y. – Antioxidant and compositional HPLC analysis of three common bamboo leaves, *Molecules* 25 (2020) 409. <https://doi.org/10.3390/molecules25020409>
18. Yu, Y., Li, Z., Cao, G., Huang, S., Yang, H. – Bamboo leaf flavonoids extracts alleviate oxidative stress in HepG2 cells via naturally modulating reactive oxygen species production and Nrf2-mediated antioxidant defense responses, *J. Food Sci.* 84 (6) (2019) 1609–1618. <https://doi.org/10.1111/1750-3841.14609>
19. Choi, M.H., Yang, S.H., & Shin, H.J. – Inhibition of melanogenesis by domestic bamboo leaves (*Sasa coreana* Nakai) extract in B16F10 melanoma cells, *Korean Society for Biotechnology and Bioengineering Journal* 33 (1) (2018) 41–47. <https://doi.org/10.7841/ksbbj.2018.33.1.41>
20. Tripathi, Y.C., Jhumka, Z., & Anjum, N. – Evaluation of total polyphenol and antioxidant activity of leaves of *Bambusa nutans* and *Bambusa vulgaris*, *Journal of Pharmacy Research* 9 (4) (2015) 271–277
21. Gong, J., Xia, D., Huang, J., Ge, Q., Mao, J., Liu, S., Zhang, Y. – Functional components of bamboo shavings and bamboo leaf extracts and their antioxidant activities *in vitro*, *J. Med. Food* 17 (11) (2014) 1208–1216. <https://doi.org/10.1089/jmf.2014.3189>
22. Tanaka, A., Zhu, Q., Tan, H., Horiba, H., Ohnuki, K., Mori, Y., Yamauchi, R., Ishikawa, H., Iwamoto, A., Kawahara, H., Shimizu, K. – Biological activities and phytochemical profiles of extracts from different parts of bamboo (*Phyllostachys pubescens*), *Molecules* 19 (6) (2014) 8238–8260. <https://doi.org/10.3390/molecules19068238>

23. Ni, Q., Xu, G., Wang, Z., Gao, Q., Wang, S., Zhang, Y. – Seasonal variations of the antioxidant composition in ground bamboo *Sasa argenteostriatus* leaves, *Int. J. Mol. Sci.* 13 (2) (2012) 2249–2262. <https://doi.org/10.3390/ijms13022249>
24. Lu, B., Wu, X., Shi, J., Dong, Y., & Zhang, Y. – Toxicology and safety of antioxidant of bamboo leaves. Part 2: Developmental toxicity test in rats with antioxidant of bamboo leaves, *Food and Chemical Toxicology* 44 (2006) 1739–1743
25. Panee, J. – Potential medicinal application and toxicity evaluation of extracts from bamboo plants, *Cellular and Molecular Biology* 9 (23) (2015) 681–692
26. Wu, D., Chen, J., Lu, B., Xiong, L., He, Y., Zhang, Y. – Application of near infrared spectroscopy for the rapid determination of antioxidant activity of bamboo leaf extract, *Food Chem.* 135 (4) (2012) 2147–2156. <https://doi.org/10.1016/j.foodchem.2012.07.011>
27. Liu, H.-C., Hsieh, C.-Y., Tsai, P.-W., Chou, T.-Y., Yang, S.-C., Chang, C.-H., Huang, Y.-P., Chien, C.-C., Lee, S.-C., Shih, H.-D., Liu, Y.-C. – Development and applications of *Bambusa stenostachya* leaf extract in personal care products, *Processes* 13 (1) (2025) 233. <https://doi.org/10.3390/pr13010233>
28. Ni, Q., Zhang, Y., Xu, G., Gao, Q., Gong, L., & Zhang, Y. – Influence of harvest season and drying method on the antioxidant activity and active compounds of two bamboo grass leaves, *Journal of Food Processing and Preservation* 37 (2013) 639–647. <https://doi.org/10.1111/jfpp.12116>
29. Zhou, Y., Chen, M., Huo, X., Xu, Q., Wu, L., & Wang, L. – Separation of flavonoids and purification of chlorogenic acid from bamboo leaves extraction residues by combination of macroporous resin and high-speed counter-current chromatography, *Molecules* 28, 4443 (2023). <https://doi.org/10.3390/molecules28114443>
30. Li, X., Tao, W., Xun, H., Yao, X., Wang, J., Sun, J., Yue, Y., Tang, F. – Simultaneous determination of flavonoids from bamboo leaf extracts using liquid chromatography-tandem mass spectrometry, *Rev. Bras. Farmacogn.* 31 (2021) 347–352. <https://doi.org/10.1007/s43450-021-00158-1>
31. Shao, S.-Y., Ting, Y., Wang, J., Sun, J., & Guo, X.-F. – Characterization and Identification of the Major Flavonoids in *Phyllostachys edulis* Leaf Extract by UPLC–QTOF–MS/MS, *Acta Chromatographica* 32(4) (2020) 228–237. <https://doi.org/10.1556/1326.2019.00688>
32. Tang, Y., Wei, X., Zou, Y., Chen, Y., & Ma, C. – Study on the Extraction Mechanism of Bamboo Flavonoids by Ultrasonic-Microwave Synergistic Extraction, *Botanical Research* 8(3) (2019) 247–255. <https://doi.org/10.12677/br.2019.83033>
33. Wedler, J., Daubitz, T., Schlotterbeck, G., & Butterweck, V. – *In vitro* anti-inflammatory and wound-healing potential of a *Phyllostachys edulis* leaf extract – Identification of isoorientin as an active compound, *Planta Medica*, 80 (2014), 1678–1684. <https://doi.org/10.1055/s-0034-1383195>
34. Lv, Z.-L., Lin, X., Miao, Z.-H., Guo, H.-X., Wang, J.-A.-H., Lei, M.-L., Pan, Y., & Zhang, B.-L. – Antioxidant activity of bamboo-leaf extracts from the species *Dendrocalamopsis oldhami*, *Scientific Research and Essays* 7(44) (2012), 3789–3796. <https://doi.org/10.5897/SRE11.1864>
35. Lu, B., Wu, X., Tie, X., Zhang, Y., Zhang, Y. – Toxicology and safety of anti-oxidant of bamboo leaves. Part 1: acute and subchronic toxicity studies on anti-oxidant of bamboo leaves, *Food Chem. Toxicol* 43 (2005) 783–792. <https://doi.org/10.1016/j.fct.2005.01.019>

36. Zhao, Y., Zhao, Y.-Y., Du, Y., Kang, J.S. – Characterization and classification of three common Bambusoideae species in Korea by an HPLC-based analytical platform coupled with multivariate statistical analysis, *Ind. Crops Prod.* 130 (2019) 389–397. <https://doi.org/10.1016/j.indcrop.2018.12.096>
37. Shen, D.-Y., Song, H.-L., Zou, T.-T., Wan, S.-Y., & Li, M.-K. – Characterization of odor-active compounds in moso bamboo (*Phyllostachys pubescens* Mazel) leaf via gas chromatography-ion mobility spectrometry, one- and two-dimensional gas chromatography-olfactory-mass spectrometry, and electronic nose, *Food Research International* 152 (2022) 110916. <https://doi.org/10.1016/j.foodres.2021.110916>
38. Jin, Y.-C., Yuan, K., Zhang, J. – Chemical composition, and antioxidant and antimicrobial activities of essential oil of *Phyllostachys heterocycla* cv. *pubescens* varieties from China, *Molecules* 16 (6) (2011) 4318–4327. <https://doi.org/10.3390/molecules16054318>
39. Wang, H., Su, Z., Deng, P., Chen, L., Yang, M., Xu, X. – Characterization of volatile organic compounds and aroma of eight bamboo species leaves, *Horticulturae* 10 (4) (2024) 394. <https://doi.org/10.3390/horticulturae10040394>
40. Kim, J.-S., Lee, H.C., Jo, J.-S., Jung, J.Y., Ha, Y.L., Yang, J.-K. – Evaluation of antioxidant and anticancer activity of steam extract from the bamboo species, *J. Korean Wood Sci. Technol.* 42 (5) (2014) 543–554. <https://doi.org/10.5658/WOOD.2014.42.5.543>
41. Menchavez, M.T., Catipay, J.D., Espra, A.S., Castillo, P.M. – Antibacterial properties of *Bambusa vulgaris* (bamboo) leaves and *Eryngium foetidum* (culantro) leaves against *Staphylococcus aureus* and *Escherichia coli* bacteria, *Glob. J. Med. Plant Res.* 6 (1) (2018) 6–13. <https://doi.org/10.22587/gjmpr.2018.6.1.2>
42. Sriraman, S., Ramanujam, G.M., Ramasamy, M., Dubey, G.P. – Identification of beta-sitosterol and stigmasterol in *Bambusa bambos* (L.) Voss leaf extract using HPLC and its estrogenic effect in vitro, *J. Pharm. Biomed. Anal.* 117 (2016) 244–250. <https://doi.org/10.1016/j.jpba.2015.06.024>
43. Ji Hyeon Kwon, Sung Yeoun Hwang, Ji Sook Han – Bamboo (*Phyllostachys bambusoides*) leaf extracts inhibit adipogenesis by regulating adipogenic transcription factors and enzymes in 3T3-L1 adipocytes, *Food Science and Biotechnology* 26 (4) (2017) 1037–1044. <https://doi.org/10.1007/s10068-017-0150-y>
44. Mao, J.-W., Yin, J., Ge, Q., Jiang, Z.-L., Gong, J.-Y. – In vitro antioxidant activities of polysaccharides extracted from Moso Bamboo-Leaf, *Int. J. Biol. Macromol.* 55 (2013) 1–5. <https://doi.org/10.1016/j.ijbiomac.2012.12.027>
45. Ma, Y., Yang, Y., Gao, J., Feng, J., Shang, Y., Wei, Z. – Phenolics and antioxidant activity of bamboo leaves soup as affected by in vitro digestion, *Food Chem. Toxicol.* 135 (2020) 110941. <https://doi.org/10.1016/j.fct.2019.110941>
46. Xiao, Z., Zhang, Q., Dai, J., Wang, X., Yang, Q., Cai, C., Mao, J., Ge, Q. – Structural characterization, antioxidant and antimicrobial activity of water-soluble polysaccharides from bamboo (*Phyllostachys pubescens* Mazel) leaves, *Int. J. Biol. Macromol.* 146 (2020) 985–993. <https://doi.org/10.1016/j.ijbiomac.2019.09.115>
47. Zulkifli, M.Z.A., Benjamin, M.A.Z., Rosdan, M.D.E.M., Saini, A., Rusdi, N.A., Awang, M.A. – Optimisation of yield, flavonoids, and antioxidant activity via ultrasound-assisted extraction of bamboo leaves from *Dinochloa sublaevigata* S. Dransf. (Wadan) in Sabah, Malaysia, *Adv. Bamboo Sci.* 10 (2025) 100128. <https://doi.org/10.1016/j.bamboo.2025.100128>

48. Jayarambabu, N., Velupla, S., Akshaykranth, A., Anitha, N., Rao, T.V. – Bambusa arundinacea leaves extract-derived Ag NPs: evaluation of the photocatalytic, antioxidant, antibacterial, and anticancer activities, *Appl. Phys. A* 129 (2023) 13. <https://doi.org/10.1007/s00339-022-06279-1>
49. Singhal, P. – Bamboo Leaves: An Emerging Multi-functional Food, *Food and Drug Safety*, 1(1) (2023) 1–9. <https://doi.org/10.55121/fds.v1i1.134>
50. Elengoe, A., Tan, C.H., Poddar, S. – Computational analysis of antimicrobial phytochemicals from *Bambusa vulgaris* Schrad. ex J.C.Wendl.: Pharmacokinetics, toxicity, and molecular docking, *Adv. Bamboo Sci.* 11 (2025) 100157. <https://doi.org/10.1016/j.bamboo.2025.100157>
51. Kalyan, N., Indira, A., Joshi, B., Rani, M., Lathwal, M., Kuhad, A., Chongtham, N. – An overview of antimicrobial potential of bamboo, *Proc. 12th World Bamboo Congress, Taiwan* (2024) 18–22 April
52. Liao, M., Ren, X., Gao, Q., Liu, N., Tang, F., Wang, G., & Cao, H. – Anti-fungal activity of moso bamboo (*Phyllostachys pubescens*) leaf extract and its development into a botanical fungicide to control pepper phytophthora blight, *Scientific Reports* 11 (2021), 14146. <https://doi.org/10.1038/s41598-021-83598-y>
53. Luis Carlos Chitiva, Mary Andrea Santamaría-Torres, Paula Rezende-Teixeira, Jessica R.P.O. Borlot, Rodrigo A. Romagna, Ximena Londono, Rodrigo R. Kitagawa, Leticia V. Costa-Lotufu, Juliet A. Prieto-Rodríguez, Ian Castro-Gamboa, Geison Modesti Costa – Uncovering metabolic alterations in HCT-116 colon cancer cells upon exposure to bamboo leaf extract obtained from *Guadua incana* Londono, *Molecules* 29 (13) (2024) 2985. <https://doi.org/10.3390/molecules29132985>
54. Lee, W., Ku, S.-K., Bae, J.-S. – Vascular barrier protective effects of orientin and isoorientin in LPS-induced inflammation in vitro and in vivo, *Vasc. Pharmacol.* 62 (1) (2014) 3–14. <https://doi.org/10.1016/j.vph.2014.04.006>
55. Gurjar, V.S., Sharma, Y.K. – Evaluation of anti-diabetic activity of *Bambusa bambos* using streptozocin diabetic model induced, *J. Biomed. Pharm. Res.* 11 (2) (2022) 44–51. <https://doi.org/10.32553/jbpr.v11i2.909>
56. Liu, J.-X., Zhu, M.-Y., Feng, C.-Y., Ding, H.-B., Zhan, Y., Zhao, Z., Ding, Y.-M. – Bamboo leaf extract improves spatial learning ability in a rat model with senile dementia, *J. Zhejiang Univ-Sci B (Biomed. & Biotechnol.)* 16 (7) (2015) 593–601. <https://doi.org/10.1631/jzus.B1400249>
57. Kim, B., Lee, K.Y., Park, B. – Isoorientin Inhibits Amyloid β_{25-35} -Induced Neuronal Inflammation in BV2 Cells by Blocking the NF- κ B Signaling Pathway, *Molecules* 26 (2021) 7056. <https://doi.org/10.3390/molecules26227056>
58. Dong Ho SHIN, Pan Gu JANG, Ki Seok OH, Jae Ha KIM, Hee Jong CHUNG and Jong Choon KIM – A Single Oral Dose Toxicity Study of Bamboo Leaf Water Extract in Sprague-Dawley Rats, *The Journal of Applied Pharmacology* 11, 200–203 (2003).
59. Funmilayo Abimbola OKENIYI, Bolanle Moji ADENIRAN, Razaq Adekunle ANIMASHAHUN, Reuben Chukwuka OKOCHA, Samuel Oyewale OLAWOYE, Olayinka Olubunmi ALABI, Oyegunle Emmanuel OKE, Ayoola John SHOYOMBO, Emmanuel Oladiran BABARINDE – Antioxidant status, hematology, performance, organ and carcass evaluation of heat-stressed broiler chickens fed with dietary *Bambusa vulgaris* leaf meal, *Veterinary Research Notes* 3 (4) (2023) 29–35. <http://doi.org/10.5455/vrn.2023.c25>

60. Gabriel Adedotun WILLIAMS, Wasiru Ajani OLAYEMI, Idowu Aminat OYEKAN, Similoluwa Aramide MUIBI, Abimbola Oladele OSO – Oral administration of aqueous bamboo leaf extract: effect on performance, haematological indices and blood oxidative status of broilers, *Agricultura Tropica et Subtropica* 57 (OV) (2024) 171–181. <https://doi.org/10.2478/ats-2024-0017>
61. Guadua Bamboo – Bamboo genera around the world, *Guadua Bamboo* (2023). Available from: <https://www.guadubamboo.com/blog/bamboo-genera> [Accessed 18 Apr. 2025]
62. Susan CANAVAN, David M. RICHARDSON, Vernon VISSER, Johannes J. Le ROUX, Maria S. VORONTSOVA, John R. U. WILSON – The global distribution of bamboos: assessing correlates of introduction and invasion, *AoB PLANTS* 9 (1) (2017) plw078. <https://doi.org/10.1093/aobpla/plw078>
63. Food and Agriculture Organization of the United Nations – Global Forest Resources Assessment 2020: Main report, FAO (2020). Available from: <https://www.fao.org/forest-monitoring/news-and-events/news/news-detail/Putting-bamboo-on-the-map/en> [Accessed 18 Apr. 2025]
64. Kiran PAUDYAL, Li YANXIA, Trinh Thang LONG, Shankar ADHIKARI, Soni LAMA, Kishor Prasad BHATTA – Ecosystem services from bamboo forests: key findings, lessons learnt and call for actions from global synthesis, INBAR Working Paper, International Bamboo and Rattan Organisation (2022)
65. Julian EVANS – Plantation Forestry in the Tropics, 3rd ed., Oxford University Press, Oxford (2009)
66. World Agroforestry Centre (ICRAF) – Annual Report 2020, World Agroforestry Centre, Nairobi (2020)
67. Business Wire – The Global Bamboo Market is Projected to Reach \$82.90 Billion by 2028 at a CAGR of 5.7% – ResearchAndMarkets.com (2021 Nov 30). Available from: <https://www.businesswire.com/news/home/20211130005687/en/The-Global-Bamboo-Market-is-Projected-to-Rreach-%2482.90-Billion-by-2028-at-a-CAGR-of-5.7---ResearchAndMarkets.com> [Accessed 18 Apr. 2025]
68. Yannick KUEHL – Chapter 4: Resources, yield, and volume of bamboos, in: W. Liese, M. Köhl (eds.), *Bamboo, Tropical Forestry* 10, Springer International Publishing, Switzerland (2015). https://doi.org/10.1007/978-3-319-14133-6_4
69. Vuong Thi THUY, Nguyen Thi HUYEN, Le Hoang TU, Nguyen Kim LOI – Status of bamboos in Binh Duong province, Vietnam: Distribution, species diversity, conservation and utilization, *Trees, Forests and People* 6 (2021) 100137. <https://doi.org/10.1016/j.tfp.2021.100137>
70. Zishan AHMAD, Anamica UPADHYAY, Yulong DING, Abolghassem EMAMVERDIAN, Anwar SHAHZAD – Chapter 1: Bamboo: Origin, habitat, distributions and global prospective, in: Z. Ahmad et al. (eds.), *Biotechnological Advances in Bamboo*, Springer Nature Singapore (2021). https://doi.org/10.1007/978-981-16-1310-4_1
71. Nguyen Hoang NGHIA, Tran Van TIEN – Bamboo Resources in Vietnam, Forest Science Institute of Vietnam (2009). Available from: <https://www.researchgate.net/publication/341071902> [Accessed 18 Apr. 2025]
72. Chu KHOI – Vietnam's bamboo sector facing the challenges of integration, *VnEconomy – Vietnam Economic Times*, Hanoi (2022 Aug 19). Available from:

- <https://vneconomy.vn/nganh-tre-viet-nam-truoc-thach-thuc-hoi-nhap.htm> [Accessed 18 Apr. 2025]
73. Tran Van HIEP – Key factors affecting small bamboo enterprises upgrading in North Vietnam: Case studies from Chuong My, Hanoi and Thanh Hoa province, Doctoral dissertation, Technische Universität Dresden, Germany (2021)
 74. Nguyen Hoang NGHIA, Nguyen Van THO, Nguyen VIEN, Pham Quang TIEN, Le Thi Mai LINH, Nguyen Thi Hong MAI – Genetic diversity assessment of two bamboo species belonging to the genus *Dendrocalamus* (Nees) in northern Vietnam based on ISSR molecular markers, *Journal of Vietnamese Forestry Science*, No. 1 (2018) 17–26. Available from: <https://www.researchgate.net/publication/336362676> [Accessed 18 Apr. 2025]
 75. Tran Van TIEN – A synopsis of the subtribe Bambusinae (Gramineae: Bambusoideae) in Vietnam, *Dalat University Journal of Science* 6 (4) (2016) 442–449.
 76. Pham Hoang HO – Flora of Vietnam: An Illustrated Flora of Vietnam, Volume III, Tre Publishing House, Ho Chi Minh City (2003), 167 pages
 77. Vinh, T.T. – Study on the Diversity of the Bamboo Subfamily (*Bambusoideae*) in the Central Highlands. PhD Dissertation, Graduate University of Science and Technology, Vietnam Academy of Science and Technology, Lam Dong, Vietnam, 2024 (in Vietnamese)
 78. Thai Vinh T., Van Duy N., Thanh Truong H., Van Tien T. (2023) – *Yersinochloa nghiana*, a new species (Poaceae, *Bambusoideae*, *Bambuseae*) from southern Vietnam. *PhytoKeys* 224, 175–182. <https://doi.org/10.3897/phytokeys.224.101201>
 79. Tran Thai Vinh, Hoang Nghia Nguyen, Nong Van Duy and Tran Van Tien – *Chimonocalamus bidoupensis*, a new temperate bamboo species (Poaceae: Bambusoideae) from Bidoup National Park, southern Vietnam, *Nordic Journal of Botany* (2021) e02853. <https://doi.org/10.1111/njb.02853>
 80. Tran Van Tien, Nguyen Hoang Nghia and Xia Nian-He – *Annamocalamus*, a new genus of bamboo (Poaceae) from Vietnam, *Candollea* 68 (1) (2013) 159–165. <https://doi.org/10.15553/c2013v681a23>
 81. Hoang Nghia Nguyen and Van Tien Tran – *Nianhochloa* gen. nov. (Poaceae, *Bambusoideae*), a new bamboo genus endemic to Bidoup Mountain, southern Vietnam, *Adansonia*, sér. 3 34 (2) (2012) 257–264. <https://doi.org/10.5252/a2012n2a5>
 82. Diep Thi My Hanh, Nong Van Duy, Tran Van Tien – Study on the applicability of bamboo in forest restoration and climate change adaptation during the 2008–2010 period, *Báo Khoa học Phổ thông* [Science Popularization Bulletin] (2010). Available from: <https://khoahocphothong.vn/vai-tro-cua-cay-tre-voi-su-ung-pho-bien-doi-khi-hau-215863.html> [Accessed 18 Apr. 2025]
 83. Department of Science and Technology – Ministry of Natural Resources and Environment – Bamboo and its role in combating climate change, Department of Science and Technology – Ministry of Natural Resources and Environment, Hanoi, Vietnam (2019). Available from: <https://vukhcn.monre.gov.vn/tin-khoa-hoc-cong-nghe-trong-nuoc/cay-tre-voi-cong-cuoc-chong-voi-bien-doi-khi-hau/tt-754> [Accessed 18 Apr. 2025]
 84. Nhan Dan Newspaper – World bamboo workshop held in Binh Duong, Nhan Dan Online Newspaper, Hanoi, Vietnam (2022 Sep 17). Available from: <https://nhandan.vn/hoi-thao-the-gioi-ve-tre-tai-binh-duong-post715690.html> [Accessed 18 Apr. 2025]

85. Andriarimalala, J.H., Kpomasse, C.C., Salgado, P., Ralisoa, N., Durai, J. – Nutritional potential of bamboo leaves for feeding dairy cattle, *Pesqui. Agropecu. Trop.* 49 (2019) e54370. <https://doi.org/10.1590/1983-40632019v4954370>
86. Utsaphong Uprarawanna, Jirapath Kaewsritong and Khongsak Srikaeo – Bamboo: Global occurrence and its significance as food and related products, *Crops* 5 (2) (2025) 11. <https://doi.org/10.3390/crops5020011>

TÓM TẮT

TỔNG QUAN TOÀN DIỆN VỀ LÁ TRE: THÀNH PHẦN HÓA HỌC, HOẠT TÍNH SINH HỌC, ĐỘC TÍNH VÀ TIỀM NĂNG TÀI NGUYÊN TRE TẠI VIỆT NAM

Lê Quỳnh Anh^{1,2,3}, Dương Thanh Tài¹, Nguyễn Thị Thu Thủy¹,
Nguyễn Cao Ngọc Hân¹, Nguyễn Thị Lan Phi^{2,3*}

¹Trường Đại học Công Thương Thành phố Hồ Chí Minh (HUIT)

²Trường Đại học Bách Khoa (HCMUT)

³Trường Đại học Quốc gia Thành phố Hồ Chí Minh (VNU HCM)

*Email: lanphi@hcmut.edu.vn

Lá tre (*Folium bambusae*) từ lâu đã được biết đến là một dược liệu truyền thống trong y học cổ truyền phương Đông, và được sử dụng phổ biến trong nhiều bài thuốc dân gian nhờ các đặc tính thanh nhiệt, lợi tiểu và chống viêm. Những năm gần đây, lá tre được nhiều nhà khoa học quan tâm, từ đó nhiều công trình nghiên cứu đã được công bố. Trong vòng hai thập kỷ qua, các nghiên cứu liên quan đến định danh loài, thành phần hóa học, hoạt tính sinh học và độc tính của lá tre đã được nghiên cứu và đăng bài liên tục. Bài báo tổng quan này nhằm hệ thống hóa và tóm lược các nghiên cứu khoa học hiện có về thành phần hóa học, hoạt tính sinh học và độc tính của chiết xuất lá tre. Nguồn dữ liệu được thu thập từ các cơ sở dữ liệu khoa học quốc tế uy tín như PubMed, ScienceDirect, Google Scholar, Web of Science, Scopus và ScienceGate, cùng với một số nguồn trực tuyến đáng tin cậy khác. Ngoài việc khảo sát hệ thống phân loại tre và hiện trạng tài nguyên tre trên phạm vi toàn cầu, bài báo đặc biệt nhấn mạnh đến thực trạng và tiềm năng phát triển tài nguyên tre tại Việt Nam, đồng thời xem xét vai trò tiềm năng của lá tre trong ngành công nghiệp thực phẩm. Tổng cộng 86 tài liệu tham khảo đã được tuyển chọn kỹ lưỡng dựa trên tiêu chí mức độ liên quan và độ tin cậy khoa học, được sử dụng để trích xuất và tổng hợp dữ liệu. Bài viết cung cấp một cái nhìn toàn diện về nguồn tài nguyên tre và các khía cạnh nghiên cứu liên quan đến lá tre về Phytochemical, hoạt tính sinh học và thử nghiệm độc tính; qua đó định hướng cho các nghiên cứu chuyên sâu hơn cũng như các thử nghiệm ứng dụng rộng rãi trong tương lai. Những đóng góp này kỳ vọng sẽ nâng cao giá trị của cây tre, thúc đẩy sự phát triển bền vững hệ sinh thái tre, và góp phần vào mục tiêu phát triển xanh.

Từ khóa: Lá tre, hóa thực vật, hoạt tính sinh học, độc tính, tài nguyên tre.